

Sea Ice Outlook
2017 June Report
Individual Outlook

Name of Contributor of Name of Contributing Organization:

CNRM (Chevallier et al)

Is this contribution from a person or group not affiliated with a research organization?

Name and organization for all contributors. Indicate primary contact and total number of people who may have contributed to your Outlook, even if not included on the author list.

Matthieu Chevallier, Constantin Ardilouze, Lauriane Batté (CNRM, Meteo France, Toulouse, France)

Do you want your June contribution to be automatically included in subsequent reports? (If yes, you may still update your contribution via a form like this one.)

No do not use my prediction this month in later months

What is the type of you outlook projection?

Dynamic Model

Starting in 2017 we are accepting both pan-Arctic and pan-Antarctic sea ice extent (either one or both) of the September monthly mean. As in 2016, we are also collecting Alaskan regional sea ice extent. To be consistent with the validating sea ice extent index from NSIDC, if possible, please first compute the average sea ice concentration for the month and then compute the extent as the sum of cell areas > 15%.

a) Pan-Arctic September extent prediction in million square kilometers.

4.39 (STD 0.60)

b) same as in (a) but for pan-Antarctic. If your method differs substantially from that for the Arctic, please enter it as a separate submission.

17.80 (STD 0.31)

c) same as in (b) but for the Alaskan region. Please also tell us the maximum possible extent if every ocean cell in your region were ice covered.

"Executive summary" of your Outlook contribution (using 300 words or less) describe how and why your contribution was formulated. To the extent possible, use non-technical language.

Sea Ice Outlook
2017 June Report
Individual Outlook

CNRM outlook is based on the operational seasonal forecast issued by Météo France in early June 2017 with the System 5 (component of the European multi-model EUROSIP).

Brief explanation of Outlook method (using 300 words or less).

The CNRM outlook is a model estimate based on a dynamical ensemble forecast. Initial conditions from the week before 1 June 2017 are used. We generate a total 51 members by adding statistical perturbations during the run.

Tell us the dataset used for your initial Sea Ice Concentration (SIC). Include name and date (e.g., "NASA Team, May 2017"). We also encourage you to submit initial fields to the dropbox, see <https://www.arcus.org/sipn/sea-ice-outlook/2017/june/call> in the section on "Submitting Figures and Gridded Data of Full Spatial Fields (Optional) of Forecasts and Initial Conditions" for detailed instructions. Required if sea Ice concentration is used.

Initial conditions for the ocean and sea ice (concentration and thickness) are provided by Mercator Océan. Basis is the Mercator Océan operational analysis, run at a 1/4° horizontal resolution using NEMO-LIM2 and the SAM ocean data assimilation system. There is no data assimilation of sea ice concentration in this analysis. The 1/4° analysis is upscaled to the 1° horizontal grid of CNRM-CM. These fields are used to nudge the ocean-sea ice component of CNRM-CM (NEMO3.2-Gelato5, 1° resolution), run in forced mode (forced by ECMWF Op. analysis). A strong restoring is applied towards Mercator SST, which acts as a constraint on sea ice concentration. Sea ice fields (concentration, thickness...) from this 1° run are used to initialize CNRM-CM (as well as ocean fields from this run).

Dataset of initial Sea Ice Thickness (SIT) used (include name and date):

Sea ice thickness information is output from the 1° simulation described above.

If you use a dynamic model, please specify the name of the model as a whole and each component including version numbers and how the component is initialized:

Météo France System 5
Component Name Initialization
Atmosphere ARPEGE-Climat v6 ECMWF Operational Analysis
Ocean NEMO3.2 Based on Mercator (DA SEEK)
Sea ice GELATO 5 Based on Mercator (no DA)

Basis of System 5 is the global coupled model CNRM-CM5 (Voldoire et al., 2014).

Horizontal resolution of the atmosphere component is T255 (70km). Nominal resolution of the ocean-sea ice component is 1° at the equator (nearly 50km in the Arctic). The sea ice model uses 4 sea ice categories (0-0.3; 0.3-0.8, 0.8-3 and >3m).

Sea Ice Outlook
2017 June Report
Individual Outlook

If available from your method for pan-Arctic extent prediction, please provide:

a) Uncertainty/probability estimate such as median, ranges, and/or standard deviations (specify what you are providing).

For pan-Arctic SIO

Median : 4.50

25% : 3.99

75% : 4.90

Min : 3.17

Max : 5.34

For pan-Antarctic SIO

Median : 17.83

25% : 17.62

75% : 17.97

Min : 16.94

Max : 18.52

Statistics are based on the 51-member ensemble.

b) Brief explanation/assessment of basis for the uncertainty estimate (1-2 sentences).

Standard deviation of the ensemble is 0.60 million km² for Arctic, 0.31 million km² for Antarctic.

c) same as in (b) but for the Alaskan region. Please also tell us the maximum possible extent if every ocean cell in your region were ice covered. See <https://www.arcus.org/sipn/sea-ice-outlook/2017/june/call> in the section on "Instructions for Submitting an Alaskan Regional Outlook" for detailed instructions.

Post-processing includes bias correction and correction of the trend based on the hindcast (1993-2014).

d) Raw (and/or post processed) forecasts for this year and retrospective forecasts in an excel spreadsheet with one year on each row and ensemble member number on columns (specifying whether raw or post processed).

Post-processing includes bias correction and correction of the trend based on the hindcast (1993-2014).