

Sea Ice Outlook
2020 August Report
Individual Outlook

Name of contributor or name of contributing organization:

CPOM

Is this contribution from a person or group not affiliated with a research organization?

Name and organization for all contributors. Indicate primary contact and total number of people who may have contributed to your Outlook, even if not included on the author list.

David Schroeder¹, Danny Feltham¹, Daniela Flocco¹, Michael Tsamados¹

¹ CPOM, Department of Meteorology, University of Reading, U.K.

² CPOM, UCL, London, U.K.

**Do you want your June contribution to automatically be included in subsequent reports?
(If yes, you may still update your contribution via the submission form.)**

Include this submission in the July report only.

What is the type of your Outlook projection?

Statistical

Starting in 2017 we are accepting both pan-Arctic and pan-Antarctic sea ice extent (either one or both) of the September monthly mean. As in 2016, we are also collecting Alaskan regional sea ice extent. To be consistent with the validating sea ice extent index from NSIDC, if possible, please first compute the average sea ice concentration for the month and then compute the extent as the sum of cell areas > 15%.

a) Pan-Arctic September extent prediction in million square kilometers.

4.3

b) same as in (a) but for pan-Antarctic. If your method differs substantially from that for the Arctic, please enter it as a separate submission.

c) same as in (b) but for the Alaskan region. Please also tell us maximum possible extent if every ocean cell in your region were ice covered.

"Executive summary" of your Outlook contribution (using 300 words or less) describe how and why your contribution was formulated. To the extent possible, use non-technical language.

We predict the September ice extent 2020 to be 3.8 (3.3-4.3) million km². This is the lowest prediction we have made based on spring melt pond fraction. The likelihood is around 30% that this September extent will be a new minimum record. In our model simulation since 1979, May 2020 has the highest mean melt pond fraction for May including some unprecedented melt pond formation in the Central Arctic during 15-18 May when air temperature exceeded 0deg C.

Brief explanation of Outlook method (using 300 words or less).

This is a statistical prediction based on the correlation between the ice area covered by melt-ponds in May and ice extent in September. The melt pond area is derived from a simulation with the sea ice model CICE in which we incorporated a physically based melt-pond model¹. See our publication in Nature Climate Change <http://www.nature.com/nclimate/journal/v4/n5/full/nclimate2203.html> for details².

References:

1. Flocco, D., Schröder, D., Feltham, D. L. & Hunke, E. C., 2012: Impact of melt ponds on Arctic sea ice simulations from 1990 to 2007. J. Geophys. Res. 117, C09032.
2. Schröder, D., D. L. Feltham, D. Flocco, M. Tsamados, 2014: September Arctic sea-ice minimum predicted by spring melt-pond fraction. Nature Clim. Change 4, 353-357, DOI: 10.1038/NCLIMATE2203.

Tell us the dataset used for your initial Sea Ice Concentration (SIC).

None.

Tell us the dataset used for your initial Sea Ice Thickness (SIT) used. Include name and date.

None.

If you use a dynamic model, please specify the name of the model as a whole and each component including version numbers and how the component is initialized:

[DynamicModelType]

If available from your method.

a) Uncertainty/probability estimates:

Median

Ranges

Standard Deviations

0.5

b) Brief explanation/assessment of basis for the uncertainty estimate (1-2 sentences).

Mean forecast error based on forecasts for the years 1984 to 2019.

c) Brief description of any post processing you have done (1-2 sentences).

See references in Section 6.